

Health Profile:


Maribor, Slovenia

*Taking cities to a
healthier future*

EURO-URHIS 2

European Urban Health Indicators System Part 2
Urban Health Monitoring and Analysis System to Inform Policy





A	Skopje, the former Yugoslav Republic of Macedonia	N	Tromsø , Norway
B	Tetovo, the former Yugoslav Republic of Macedonia	O	Bistrița, Romania
C	Bordeaux, France	P	Craiova, Romania
D	Montpellier, France	Q	Iași, Romania
E	Oberhausen, Germany	R	Bratislava, Slovakia
F	Köln, Germany	S	Košice, Slovakia
G	Liepāja, Latvia	T	Ljubljana, Slovenia
H	Riga, Latvia	U	Maribor, Slovenia
I	Kaunas, Lithuania	V	Birmingham, United Kingdom
J	Šiauliai, Lithuania	W	Cardiff, United Kingdom
K	Amsterdam, The Netherlands	X	Glasgow, United Kingdom
L	Utrecht, The Netherlands	Y	Merseyside, United Kingdom
M	Oslo, Norway	Z	Greater Manchester, United Kingdom

Depression and anxiety were less often reported in Maribor compared to the other EURO-URHIS 2 cities.

All-cause mortality in both males and females is similar in Maribor compared to other EURO-URHIS 2 cities. Mortality from malignant neoplasms and from diseases of the circulatory and respiratory system does not differ.

Heavy episodic drinking in Maribor youth occurs more often than in other EURO-URHIS 2 cities, whereas binge drinking in adults occurs less often. Smoking in both youth and adults occurs as often in Maribor as in the other cities.

The proportion of both youth and adults who are overweight or obese is higher than the overall EURO-URHIS 2 proportion.

Health and health determinants in Maribor vary considerably by age, gender and level of education.


This health profile describes the health situation and associated health determinants in Maribor compared with those observed in other European urban areas.

Maribor is one of the urban areas chosen for EURO-URHIS 2 (European Urban Health Indicator System Part 2), a project that aims to identify health problems in urban areas. The EURO-URHIS 2 project describes health and health determinants specific to urban areas in Europe, covering cities in North, East, South, and West Europe. This project may add to information that is already locally available, in that it is the first study to enable reliable comparisons of health status between different cities in Europe. Policy makers can use the information to prioritise topics for urban health policy and for interventions in an evidence-based way.

EURO-URHIS 2 gathered information by collecting data from routinely available registration data, and by conducting youth and adult surveys at the end of 2010. In total, data from 26 urban areas in Europe were available for between-city comparisons and benchmarking.

The routinely available registration data relate to the most recently available year (2006-2008). The youth survey was a school-based survey of 14-16 year olds. In Maribor, 430 students completed a valid questionnaire. The adult survey was carried out involving a representative sample of adults aged 19-64 and 65+. In Maribor, 441 19-64 year olds and 408 65+ year olds completed valid questionnaires.

More detailed information on the justification of methods and instruments that were used, as well as response rates, selection of cities and indicators, and statistical methodology, can be found on our websites: www.urhis.eu and <http://results.urhis.eu>. The websites also provide data from other participating urban areas and comparisons between specific cities can be made.



The graphs in this health profile show the health status of the urban area compared to other EURO-URHIS 2 urban areas. The whiskers represent the lowest and highest value within the EURO-URHIS 2 project on a scale of 0 to 100%. The grey bar represents the 25th, 50th, and 75th percentile. The urban area value is shown as a diamond, which is blue when the value is not statistically significantly different from the EURO-URHIS 2 mean and red when the difference is statistically significant (at the 5% level).




Figure 1. Age distribution




Figure 2. Level of education

Differences in health status may possibly be explained by age and education. Figures 1 and 2 show the age distribution and level of education in Maribor compared to the other EURO-URHIS 2 urban areas. Age did not explain any observed differences in the adult survey between Maribor and other EURO-URHIS 2 urban areas. Education differences between adults from Maribor and other EURO-URHIS 2 cities could explain the significantly different level of the sense of belonging to the immediate neighbourhood.

DISCLAIMER

To achieve maximum quality of the data, all instruments used were based on knowledge of earlier studies and expert consultations, and were piloted, validated, and optimised. The survey questionnaires of EURO-URHIS 2 were based on already existing, validated instruments; selected indicators were as little culturally sensitive as possible. Questionnaires were translated in the local language(s) and, for validation purposes, back-translated into English. Youth survey response rates were generally very high. In the adult survey, a minimum response rate of 30% was required to be included for benchmarking. Despite all our efforts, and as in any survey, the point estimates for certain health indicators in your urban area may deviate from other estimates, and may not be comparable to other local information due to differences in study methodology and indicator definitions. If you would like further information regarding the methodology, please see our websites: <http://www.urhis.eu> and <http://results.urhis.eu>.

Health-related Characteristics of Maribor

	Indicator	Maribor	Slovenia	EURO-URHIS 2 range (percentiles)					EURO-URHIS 2 mean	N
				min	25th	50th	75th	max		
Demographic	1. Population size (x1,000)	182	2,010	67	264	406	708	2,565	570	23
	2. Population density	246	100	27	1,115	2,040	2,840	4,580	1,974	24
	3. Population aged 0-19 years	17%	20%	17%	20%	22%	24%	28%	22%	23
	4. Population aged 65+ years	18%	16%	7%	11%	14%	15%	20%	14%	23
	5. Live births	41	53	39	45	52	58	75	53	24
	6. Teenage pregnancies	10	5	4	7	11	20	33	14	18
	7. Pregnancies after age 35	19	21	7	18	23	33	59	28	18
Socio-economic	8. Unemployment (age 19-64)	7.8%	-	3.6%	4.0%	4.9%	7.2%	10.2%	5.8%	16
	9. Higher level education	25%	-	25%	33%	45%	53%	72%	45%	16
	10. Not enough money	22%	-	5%	11%	16%	22%	61%	21%	16
	11. Low family wealth	5%	-	5%	7%	13%	21%	44%	16%	20
Health System	12. MMR vaccinated	-	96%	83%	88%	94%	97%	100%	93%	19
	13. DTP vaccinated	-	97%	83%	93%	95%	97%	99%	94%	19
	14. Cervical smear test	74%	-	41%	62%	70%	76%	83%	68%	16
	15. Cholesterol measurement	48%	-	23%	42%	47%	52%	64%	47%	16
Health Status	16. Life expectancy - male	-	75.5	68.2	71.0	75.3	76.1	77.0	73.6	18
	17. Life expectancy - female	-	82.6	76.2	78.5	80.2	81.0	82.0	79.7	18
	18. Infant mortality	3.0	2.4	1.3	3.5	4.9	5.7	9.4	5.0	24
	19. Low birth weight	9.0%	6.3%	2.7%	5.2%	6.6%	8.1%	11.8%	6.7%	22

Table 1. Health-related characteristics of Maribor

Source. Indicators 1-7, 12-13, and 16-19: routinely available registration data; indicators 8-10 and 14-15: adult survey; indicator 11: youth survey. Missing data are indicated by "-".

N = number of urban areas that were able to collect data on the specific indicator.

1. number of inhabitants; **2.** number of inhabitants per km²; **3.** % of inhabitants aged 0-19 years; **4.** % of inhabitants aged 65 years or older; **5.** number of births per 1,000 women aged 15-44 years; **6.** number of births per 1,000 women aged 15-19 years; **7.** number of births per 1,000 women aged 35-44 years; **8.** % of adults aged 19-64 years who are unemployed; **9.** % of adults who attained higher level education; **10.** % of adults who do not have enough money for daily expenses; **11.** % of youth who live in a low wealth family, as defined by a FAS (Family Affluence Scale) score of ≤3; **12.** % of population who have completed measles, mumps, and rubella (MMR) vaccination courses before school-age; **13.** % of population who have completed diphtheria, tetanus, and poliomyelitis (DTP) vaccination courses before school-age; **14.** % of adult women who have undergone a cervical smear test within the past three years; **15.** % of adults who had their serum cholesterol measured within the last year; **16-17.** number of years that a newborn is expected to live if current mortality rates continue to apply; **18.** annual number of deaths of children under one year of age, per 1,000 births; **19.** % of total live births weighing less than 2,500 grams

Compared to other cities in EURO-URHIS 2, Maribor is an urban area with low population density and a somewhat older population. The number of annual live births in Maribor is lower than the overall EURO-URHIS 2 mean.

The percentage of inhabitants with higher level education in Maribor (25%) is relatively low compared to the overall EURO-URHIS 2 mean. The proportion of adults who reported to not have enough money for daily expenses (22%) is significantly higher than in the other EURO-URHIS 2 urban areas. The percentage of youth that reported to live in poor families (5%) is significantly lower than the EURO-URHIS 2 mean.

Significantly more women have undergone a cervical smear test

compared to the overall EURO-URHIS 2 average.

Infant mortality is an indicator for population health and quality of health care services. With an infant mortality rate of 3.0 per 1,000 live births, Maribor belongs to the 25% of urban areas in EURO-URHIS 2 with the lowest infant mortality.

At the population level, low birth weight is an indicator for pregnancy conditions and perinatal care. Low birth weight can at the individual level also result in health problems later in life. Of all newborns in Maribor, 9.0% had a low birth weight, which is higher than the overall EURO-URHIS 2 mean.

YOUTH HEALTH STATUS

	Indicator	Maribor	EURO-URHIS 2 range (percentiles)			EURO-URHIS 2 mean	N
			0%	50%	100%		
Health Status	1. Good self-perceived health	97%				92%	20
	2. Elevated risk of psychological problems	12%				20%	20
	3. Psychosomatic symptoms	9%				10%	20
	4. Low back pain	44%				42%	20
Lifestyle Factors	5. Overweight and obesity	17%				13%	15
	6. Physical activity ≥2 hours/week	53%				50%	20
	7. Regular fruit consumption	57%				49%	20
	8. Regular vegetable/salad consumption	66%				52%	20
	9. Regular tooth brushing	77%				72%	20
	10. Frequently watching television	47%				60%	20
	11. Daily smoking	12%				12%	20
	12. First smoking ≤13 years	26%				24%	20
	13. Heavy episodic drinking	40%				33%	20
	14. First alcohol ≤13 years	67%				53%	19
	15. Ever used cannabis	14%				16%	20
	16. Unprotected sexual intercourse	3%				4%	20
Environ- ment	17. Crime in area	19%				35%	20
	18. Involved in traffic accident	8%				7%	18
	19. Being bullied	4%				7%	20

Table 2. Health status and determinants in youth (14-16 years)

Source. Indicators 1-19: youth survey. Missing data are indicated by "-".

N = number of urban areas that were able to collect data on the specific indicator.

1. % of youth who perceive their health as good, very good, or excellent; **2.** % of youth with an overall Strengths and Difficulties Questionnaire (SDQ) score of 20 or higher; **3.** % of youth who reported a lot of headaches, stomach aches, or sickness during the past six months; **4.** % of youth who experienced low back pain during the past month; **5.** % of youth overweight or obese according to the international BMI cut-offs; **6.** % of youth who participate in vigorous physical activity for more than two hours per week in their free time; **7.** % of youth who eat fruit on most days of the week; **8.** % of youth who eat vegetables and/or salads on most days of the week; **9.** % of youth who brush their teeth more than once a day; **10.** % of youth who watch television for more than two hours on weekdays; **11.** % of youth who smoke tobacco every day; **12.** % of youth who reported first smoking at ≤13 years; **13.** % of youth who drank five or more units of alcohol on one occasion during the past 30 days; **14.** % of youth who reported first drinking alcohol at ≤13 years; **15.** % of youth who ever used cannabis; **16.** % of the total youth population who did not use a condom the last time they had sexual intercourse; **17.** % of youth who reported presence of crime, violence, or vandalism in the area where they live; **18.** % of youth who had a road traffic accident resulting in injury over the past 12 months; **19.** % of youth who have been bullied at least twice in the past couple of months

Health Status and Determinants in Youth

Table 2 gives an overview of the health status and determinants in Maribor youth, as reported from the survey. Self-perceived health is a measure of adolescent well-being. 97% of youth in Maribor perceived their health to be (very) good or excellent, which is significantly higher than the overall EURO-URHIS 2 proportion. In Maribor, a significantly lower proportion of youth were identified with an elevated risk of psychological problems (12%), compared to the overall EURO-URHIS 2 proportion.

Childhood obesity is related to a higher risk of obesity, disability, and premature death later in life. In Maribor, 17% of youth are overweight or obese, which is significantly higher than the overall EURO-URHIS 2 proportion. Physical activity can contribute to maintaining a healthy weight and preventing the occurrence of chronic conditions. Furthermore, physical activity

is associated with psychological benefits and with a better school performance in young people. The proportion of youth who reported participation in vigorous physical activity for two or more hours per week is similar in Maribor (53%), compared to the overall EURO-URHIS 2 proportion. Sedentary behaviour is related to overweight and obesity, independent of physical activity. Youth in Maribor watch significantly less television on weekdays compared to other urban areas in EURO-URHIS 2. A healthy diet can lower the risk of obesity. Regular consumption of fruit and vegetables occurs more frequently in Maribor than in other EURO-URHIS 2 urban areas.

Significantly more students in Maribor brush their teeth at least twice a day compared to other EURO-URHIS 2 cities.

Initiation of smoking and drinking alcohol at a young age is a strong predictor of smoking during adulthood and of later problems with alcohol. The proportion of youth in Maribor who smoke daily (12%) is similar to the overall EURO-URHIS 2 proportion. Drinking alcohol at the age of 13 or younger occurs significantly more often in Maribor than in other EURO-URHIS 2 urban areas. Also heavy episodic drinking of five or more units of alcohol on one occasion was reported significantly more often in Maribor (40%) compared to the total EURO-URHIS 2 population.

Regular cannabis use in young people can lead to impaired cognitive development. 14% of youth in Maribor have ever used cannabis, which is similar to the overall EURO-URHIS 2 proportion.

Neighbourhood crime, violence, or vandalism was significantly less often reported by youth in Maribor (19%) compared to other cities. The proportion of youth who were victims of bullying in the past couple of months was significantly lower.

ADULT HEALTH STATUS

Indicator		Maribor	Slovenia	EURO-URHIS 2 range (percentiles)					EURO-URHIS 2 mean	N
				min	25th	50th	75th	max		
Morbidity	1. HIV/AIDS incidence - male	7	5*	2	6	8	23	71	16	19
	2. HIV/AIDS incidence - female	0	0*	0	2	6	12	16	7	19
	3. Tuberculosis incidence	11	10	5	11	17	39	153	33	22
	4. Lung cancer incidence	-	62	29	42	55	62	103	54	13
Mortality	5. All-cause mortality - male	879	920	654	752	834	1,014	1,426	919	19
	6. All-cause mortality - female	500	500	362	495	542	640	821	560	19
	7. Malignant neoplasms - male	251	277	195	230	245	258	336	250	22
	8. Malignant neoplasms - female	145	145	114	143	153	162	232	154	22
	9. Diseases of the circulatory system - male	331	321	154	227	298	456	676	353	22
	10. Diseases of the circulatory system - female	204	211	91	147	199	299	406	220	22
	11. Diseases of the respiratory system - male	60	70	32	55	62	80	158	72	22
	12. Diseases of the respiratory system - female	28	30	12	21	36	50	120	43	22
	13. Transport accidents	13	14	1	3	5	11	16	7	21
	14. Suicide and intentional harm	22	23	4	8	11	15	29	12	22

Table 3. Morbidity and mortality

Source. Indicators 1-14: routinely available registration data. Missing data are indicated by "-".

* Country level data include HIV incidence only.

N = number of urban areas that were able to collect data on the specific indicator.

1-4. Number of newly diagnosed cases with a specific disease per 100,000 persons per year; **5-6.** All-cause mortality rate per 100,000 persons per year (standardised on European population); **7-14.** Mortality rate due to a specific cause per 100,000 persons per year (standardised on European population)

Health Status and Determinants in Adults

The health status of a population can be assessed by using a number of parameters, such as those referring to acute and chronic disease, mortality, psychological well-being, and self-perceived health. Table 3 and indicators 1-8 of Table 4 show the overall health status among adults in Maribor, compared to other cities in Europe. The results show that in Maribor the incidence of tuberculosis is similar to the overall

average in all EURO-URHIS 2 urban areas, whereas the incidence of HIV/AIDS in females is lower.

All-cause mortality in both males and females is comparable to other cities. Mortality from transport accidents and from suicide and intentional harm occurs more often compared to other EURO-URHIS 2 cities.

Health Status and Determinants in Adults (continued)

	Indicator	Maribor	EURO-URHIS 2 range (percentiles)			EURO-URHIS 2 mean	N
			0%	50%	100%		
Health Status	1. (Very) good self-perceived health	51%				64%	16
	2. Psychological problems	20%				23%	16
	3. Depression/anxiety	7%				9%	16
	4. Cardiovascular disease (age 65+)	21%				18%	16
	5. Cancer	2%				2%	16
	6. Asthma or bronchitis	5%				7%	16
	7. Long-standing illness with restrictions	39%				28%	16
	8. Low back pain	61%				45%	16
Lifestyle Factors	9. Regular consumption of fruit/vegetables	61%				53%	16
	10. Regular breakfast	53%				78%	16
	11. Being physically active ≥ twice a week	44%				46%	16
	12. Overweight and obesity	60%				50%	16
	13. Daily smoking	21%				18%	16
	14. Passive smoking by non-smokers	23%				13%	16
	15. Binge drinking	13%				17%	16
	16. Cannabis last year (age 19-64)	3%				5%	16
Environment	17. Green areas suitable for recreational activities	84%				84%	16
	18. Belonging to immediate neighbourhood	67%				54%	16
	19. Social cohesion in neighbourhood	51%				52%	16
	20. Exposure to severe noise	17%				14%	16
	21. Damp spots or mould at home	26%				27%	16

Table 4. Health status and determinants in adults (19 years and older)

Source. Indicators 1-21: adult survey. Missing data are indicated by “-”.

N = number of urban areas that were able to collect data on the specific indicator.

1. % of adults who perceive their health to be good or very good; **2.** % of adults with a score of four or more on the General Health Questionnaire (GHQ); **3.** % of adults who reported to be diagnosed with or treated for anxiety or depression during the past year; **4.** % of adults aged 65 years and older who were diagnosed with or treated for heart attack, angina, or heart failure during the past year; **5.** % of adults who were diagnosed with or treated for (any kind of malignant) cancer during the past year; **6.** % of adults who were diagnosed with or treated for bronchial asthma or chronic bronchitis during the past year; **7.** % of adults who suffer from any long-standing illness, long-standing effect from injury, disability, or other long-standing condition; **8.** % of adults who had low back pain longer than one day in the past month; **9.** % of adults who eat, on average, four or more portions of fruit and/or vegetables per day; **10.** % of adults who have breakfast at least four times a week; **11.** % of adults who are physically active for at least 30 minutes twice a week or more; **12.** % of adults overweight or obese, defined as a BMI of ≥ 25 kg/m²; **13.** % of adults who smoke every day; **14.** % of non-smokers who are exposed to second-hand smoking inside their home; **15.** % of adults who drink six or more portions of alcohol on one occasion, at least once a week (men) or at least once a month (women); **16.** % of adults aged 19-64 years who used cannabis during the past year; **17.** % of adults who perceive the green areas in their neighbourhood to be suitable for active recreational activities; **18.** % of adults who feel that they belong to their immediate neighbourhood; **19.** % of adults who perceive their neighbourhood to be socially cohesive; **20.** % of adults who were exposed to severe noise from outdoors during the past 12 months; **21.** % of adults who had wet or damp spots and/or mould or mildew inside their homes (other than in basements) within the past 12 months

The proportion of people in Maribor who perceive their health to be good or very good (51%) is lower than the average in the other urban areas in EURO-URHIS 2. The percentage of adults who reported psychological problems in Maribor (20%) is comparable to other urban areas in EURO-URHIS 2. Depression and anxiety were significantly less often reported in Maribor. Long-standing illness with restrictions and low back pain, on the contrary, were more prevalent.

Several lifestyle factors and environmental determinants can affect health (Table 4, indicators 9-21). Daily smoking, for instance, increases the risk of cancer, particularly lung cancer. Smokers are also at far greater risk of developing heart disease, stroke, and emphysema. Binge drinking is associated with many health problems, which include injuries and violence, sexually transmitted diseases, alcohol dependency, liver disease, and neurological damage. The percentage of persons who smoke

daily (21%) does not differ from other EURO-URHIS 2 cities, whereas exposure to second-hand smoking inside their home was more often reported in Maribor. The proportion of adults who regularly drink more than six units of alcohol (13%) is significantly lower in Maribor compared to the overall EURO-URHIS 2 mean.

Being overweight and obese are important determinants of death worldwide. They increase the risk of cardiovascular diseases, diabetes, musculoskeletal disorders, and some cancers. In Maribor, 60% of the adults are overweight or obese, which is higher than the overall EURO-URHIS 2 proportion. Being overweight and obese are related to lack of regular physical activity. Being physically active reduces the risk of hypertension, coronary heart disease, stroke, diabetes, breast and colon cancer, depression, and the risk of injury caused by falls.

The proportion of adults in Maribor physically active more than twice a week (44%) does not differ from the total EURO-UHIS 2 proportion. Adults in Maribor more frequently eat fruit and vegetables, but a regular breakfast was significantly less often reported.

Psychological well-being may be influenced both by the availability of green spaces in the neighbourhood that are suitable for recreational activities and by aspects of social

cohesion. In Maribor, 84% perceived their green spaces to be suitable for recreational activities, which is comparable to the other cities. The percentage of adults who perceived their neighbourhood to be socially cohesive was 51%, which is similar to the overall EURO-UHIS 2 average. A significantly higher percentage of people in Maribor feel that they belong to their immediate neighbourhood. Exposure to severe noise from outdoors was also more often reported compared to the other EURO-UHIS 2 cities.

Indicator			Age		Gender		Education level	
		Total Population	19 - 64	65 +	Male	Female	Secondary level or lower	Higher level
Health Status	1. (Very) good self-perceived health	51%	59%*	23%*	53%	50%	46%*	67%*
	2. Psychological problems	20%	18%*	26%*	15%*	24%*	21%	16%
	3. Long-standing illness with restrictions	39%	33%*	63%*	38%	40%	42%*	29%*
Lifestyle Factors	4. Overweight and obesity	60%	57%*	68%*	67%*	52%*	61%	55%
	5. Daily smoking	21%	25%*	6%*	22%	20%	24%*	11%*
	6. Binge drinking	13%	12%	16%	13%	12%	12%	15%
	7. Regular consumption of fruit/vegetables	61%	61%	59%	53%*	68%*	58%	67%
	8. Being physically active \geq twice a week	44%	43%	48%	48%	41%	42%*	52%*
	9. Social cohesion in neighbourhood	51%	49%*	60%*	49%	52%	51%	50%

Table 5. Health and health determinants by demographic groups in Maribor

Source. Adult survey.

Indicators are defined in Table 4. Missing data are indicated by "-".

* Statistically significant difference between subgroups at the 5% level.

Health and Health Determinants by Demographic Groups

Health and health determinants can vary considerably as according to age, gender, and education. Table 5 subdivides a selection of important health indicators in Maribor by subgroup: respondents aged 19-64 and 65+ years, males and females, and adults who achieved secondary level education or lower and higher level education.

Respondents aged 19-64 years in Maribor more often perceived their health to be good or very good, less frequently experienced psychological problems, and were less often restricted by a long-standing illness, than is the case for older respondents. Younger respondents had a lower tendency to be overweight or obese, were more likely to be daily smokers, and less often perceived their neighbourhood as being socially cohesive. Binge drinking, fruit and vegetable consumption, and physical activity did not differ by age.

Men and women in Maribor did not differ in self-perceived

health and restrictions due to long-standing illness. Neither did the percentage of daily smoking, binge drinking, physical activity, and perceived social neighbourhood cohesion differ between sexes. Men in Maribor less frequently experienced psychological problems, had a greater tendency to be overweight or obese, and less frequently ate fruit and vegetables compared to women.

Adults in Maribor who attained secondary level education or lower less often perceived their health to be good or very good and were more often restricted by a long-standing illness than adults with higher level education. Lower educated respondents were more likely to be daily smokers and less often engaged in physical activity. The occurrence of psychological problems, the proportion of overweight or obesity, binge drinking, fruit and vegetable consumption, and perceived social neighbourhood cohesion did not differ by education level.

Healthy Life Expectancy




Figure 3. Healthy Life Expectancy

Presented here are estimates of healthy life expectancy (HLE) at ages 15 and 60 for men and women in eligible EURO-URHIS 2 urban areas. HLE was calculated first by estimating life expectancy at each age using recent 5-year averages of all-causes mortality for each urban area. From this, life expectancy was broken down into years living in good and poor perceived health, estimated using responses to the EURO-URHIS 2 adult survey question: How is your health in general?: Very good/Good/Fair/Bad/Very bad/Don't know, and the youth survey question: In general, would you say your health is..?: Excellent/Very Good/Good/Fair/Poor. Those answering very good, good or fair on the adult survey were classed as being in good perceived health, with the remainder in poor perceived health. For the youth survey, fair and poor were categorised as poor perceived health to match the scale applied to the adult survey. It was then possible to calculate the total years in good and poor perceived health and present this as a population level HLE. Full details on this process will be available in the final EURO-URHIS 2 project report, available at www.urhis.eu.

Male life expectancy in Maribor at age 15 was 58.3 years. This was 3.5 years less than the highest in the sample (Oslo, 61.8 years), and 5.5 years more than the lowest (Kaunas, 52.8 years). At this age, males were estimated to spend 49.9 years in good perceived health. This is 8.8 years less than the longest HLE (Amsterdam, 58.7 years) and 4.8 years more than the shortest (Siauliai, 45.1 years).

Male life expectancy in Maribor at age 60 was 17.7 years. This was 1.9 years less than the highest in the sample (Oslo, 19.6 years), and 2.1 years more than the lowest (Siauliai, 15.6 years). At this age, males were estimated to spend 13.8 years in good perceived health. This is 3.9 years less than the longest HLE (Oslo, 17.7 years) and 3.0 years more than the shortest (Siauliai, 10.8 years).

Female life expectancy in Maribor at age 15 was 62.8 years. This was 1.3 years less than the highest in the sample (Bordeaux, 64.1 years), and 1.4 years more than the lowest (Siauliai, 61.4 years). At this age, females were estimated to spend 54.0 years in good perceived health. This is 5.7 years less than the longest HLE (Oslo, 59.7 years) and 3.1 years more than the shortest (Kaunas, 50.9 years).

Female life expectancy in Maribor at age 60 was 20.0 years. This was 0.9 years less than the highest in the sample (Bordeaux, 20.9 years), and 0.7 years more than the lowest (Siauliai, 19.3 years). At this age, females were estimated to spend 13.9 years in good perceived health. This is 5.4 years less than the longest HLE (Montpellier, 19.3 years) and 1.8 years more than the shortest (Siauliai, 12.1 years).



Landeszentrum Gesundheit
Nordrhein-Westfalen



Beneficiaries

The University of Manchester; Municipal Health Service Utrecht; University of Liverpool; The Iuliu Hatieganu University of Medicine & Pharmacy Epidemiology Department; The Norwegian Institute of Public Health; Municipal Health Service Amsterdam; Kaunas University of Medicine; Regional Public Health and Health Promotion Centre (Slovenia); Institute of Health and Work, North Rhine-Westphalia; Slovak Public Health Association; Hacettepe University, Department of Public Health; North West Regional Health Brussels Office; Latvian Public Health Agency; South East European University; National Federation of Regional Health Observatories; Pham Ngoc Thach University of Medicine

Local EURO-URHIS 2 representative in Maribor:

Dr Olivera Stanojevic-Jerkovic
Department of Social Medicine
Regional Institute of Public Health Maribor

olivera.stanojevic@zzv-mb.si
+386 (0)2 450 01 53

EURO-URHIS 2

European Urban Health Indicators System Part 2
Urban Health Monitoring and Analysis System to Inform Policy



The research leading to these results has received funding from the European Union's Seventh Framework Programme (FP7/2007-2013) under grant agreement no 223711

© EURO-URHIS 2, August 2012

Municipal Health Service Utrecht, The Netherlands

Design by Nicola Dale

